

Pre-Op Medication Reference Sheet <u>This document is intended only for patients receiving an anesthetic for</u>

elective surgical or diagnostic procedure.

Continue up to and including the	Take up to Day of Surgery,	Consult	Discontinue prior to surgery:
Day of Surgery:	<u>DO NOT</u> take on the Day of Surgery:	Prescriber. Modifications may be required:	
Alzheimer's Medications Acid Blockers, PPIs: Esomeprazole, Omeprazole, Pantoprazole	ACE inhibitors: Anything ending in – pril <u>Note:</u> Patient may take their evening dose as prescribed.	• • • •	Metformin : Stop 48 to 72 hours prior If contrast dye to be used for procedure.
Psychotropics/Mood-Stabilizer Medications: Fluvoxamine, Paroxetine, Sertraline, Venlafaxine, Lithium, Valproate, Xanax, Ativan, Benzodiazepines, SSRI, TCAs, Bupropion, SNRI	Angiotensin II receptor blockers: Anything ending in -artan Neurologic Medication: Provigil	Aspirin : IF stents present	Aspirin : If NO Cardiac or Carotid stents present: stop 7 days prior Phentermine: Hold 4 to 7 days prior
MAOIs: See reference chart #1	Direct Renin Inhibitors: Aliskiren, Amturnide, Tekamlo, Tekturna GLP-1 Receptor Agonists: <u>(daily</u> dosing) (Semaglutide, Dulaglutide, etc)	Warfarin (Coumadin)	NSAIDs: Stop 7 days prior. Diclofenac, Ibuprofen, Piroxicam, Aleve, Naprosyn Cox-2-Selective NSAIDS: Celecoxib- May be maintained per surgeon
Anti-Parkinson Medications: Cabidopa, L-Dopa, Levodopa,	Diuretics: Dyazide, Furosemide, HCTZ, chlorthalidone	Dabigatran (Pradaxa)	Herbal Supplements: Stop 7 days prior
Anti-seizure medications: Barbiturates, Carbamazepine, Phenytoin, Sodium Valproate	Particulate antacids: Tums, Maalox Insulin, short-acting (Regular): Novolog, Humalog, Apidra, Novolin R, Humulin R	Rivaroxaban (Xarelto)	Vitamin E: Stop 7 days prior GLP-1 Receptor Agonists: <u>(weekly dosing)</u> (Semaglutide, Dulaglutide, etc) Stop 7 days prior
Beta Blockers: Anything ending in -olol	Iron supplements Gout Therapy: Colchicine	Apixaban (Eliquis)	Erection Medications: Stop 48 hrs prior. Cialis, Levitra, Viagra
Anti-arrhythmic: Amiodarone, Digoxin	Oral Diabetic agents Pulmonary Agents: Theophylline	Heparin, LMWH: Dalteparin, Enoxaparin	Naltrexone: Stop 2- 3 days prior.



Pre-Op Medication Reference Sheet This document is intended only for patients receiving an anesthetic for

elective surgical or diagnostic procedure.

Other Anti-hypertensive and Cardiac	Estrogen compounds not used for	Insulin and Insulin
Medications:	birth control purposes	pumps:
(Alpha-antagonists, Calcium-channel blockers,	Psychostimulants: (ADHD meds)	See reference chart #2
Nitrates, Vasodilators):	Methylphenidate	
Clonidine, Digoxin, Diltiazem, Hydralazine,	Pain Medications: Tramadol,	
Isosorbide, Methyldopa, Verapamil, Nifedipine,	Tapentadol, Meperidine	
Hydrazaline, Nitroglycerin Patch		

Immunosuppressant and Rheumatologic Medications: Hold if risk of renal failure and not a transplant patient Adalimumab, Allopurinol, Azathioprine, Cyclosporine, Etanercept, Infliximab, Methotrexate, Sirolimus, Tacrolimus	Pulmonary Agents: Theophylline	Suboxone, Subutex
Steroids: inhaled and/or systemic		
Anti-Virals/Antiretrovirals Medications to Control Respiratory Conditions (Asthma/COPD): All Inhalers		
Birth control pills used for pregnancy prevention		
Thyroid Medications		
Medications to control chronic pain: Ex: Oxycontin, MS Contin, Fentanyl Patch, Methadone		



Pre-Op Medication Reference Sheet This document is intended only for patients receiving an anesthetic for

elective surgical or diagnostic procedure.

Selected Herbal/Alternative Medicine -Selective MAO-A Inhibitors		
Generic Name	Comment	
Resveratrol	Found in skin of red grapes	
Berberine	Found in many herbs (e.g. goldenseal)	
Selected Herbal/Altern	native Medicine Products - Non-Selective MAO-A and MAO-B Inhibitors	
Generic Name	Comment	
Curcumin	Found in turmeric	
Harmala alkaloids	Found in tobacco, Syrian rue, passion flower, ayahausca, Tribulus terrestris	
Rhodiola Rosea	Active constituents unknown	
Selected Herbal/Altern	native Medicine Products –Selective MAO-I	
Generic Name	Comment	
Catechin	Found in tea plant, cocoa, cat's claw	
Desmethoxyyangonin	Found in kava	
Epicatechin	Found in tea plant, cocoa, cat's claw	
Fo-Ti	Active constituents unknown	
Hydroxytyrosol	Found in olive oil	
Piperine	Found in pepper	
Selected Herbal/Alternative Medicine Products-Selectivity Unknown		
Generic Name	Comment	
Myristicin	Found in nutmeg, parsley, dill	
Siberian ginseng, Yerba mate, Yohimbe	Active constituents unknown	



elective surgical or diagnostic procedure.

MAOI Medications Inhibitors	Non-selective MAO-A and MAO-B	
Generic Name	Brand Name Examples	
Isocarboxazid	Marplan	
Isoniazid	Generic only	
Linezolid	Ζγνοχ	
Phenelzine	Nardil	
Procarbazine	Matulane	
Tranylcypromine	Parnate	MAOI/Herbals
MAOI Medications	Selective MAO-B Inhibitors	
Generic Name	Brand Name Examples	
Rasagiline	Azilect	
Selegiline	Eldepryl, Emsam (patch), Zelapar	



elective surgical or diagnostic procedure.

Insulins :	Preoperative Management	
Short and rapid-acting Aspart (NovoLog) Glulisine (Apidra) Lispro (Humalog, Humalog U200) Regular insulin (Humulin R, Novolin R) Powdered Human Insulin (Afrezza)	Hold dose on the morning of the procedure.	Resume usual dose when eating.
Non-Insulin Injectable: Byetta, Victoza		
Intermediate-acting Extended zinc insulin (Ultralente) NPH (Humulin N, Novolin N) Zinc insulin (Lente)	Take half of usual dose morning of the procedure.	Resume usual dose when eating.
Long acting Detemir (Levemir) Glargine/Touejeo (Lantus)	Continue up to and including the Day of Surgery.	Resume usual dose after the procedure.
Mixed insulins	Take half (50%) of usual dose morning of the procedure.	Resume usual schedule when eating.
Subcutaneous/Basal insulin infusion pumps	Consider a discussion with patient and/or endocrinologist. In general, basal rate can be continued at usual dose, and meal time corrections can be discontinued until the patient is eating.	

elective surgical or diagnostic procedure.

Instruct patient to take Day-of-Surgery dose (as scheduled) with a small sip of water

AREA: Clinical Practice	EFFECTIVE DATE: 03/01/2024
POLICY NUMBER: VE-M-002	REVIEWED/UPDATED: 02/2014, 01/2015, 03/2016, 4/2017, 1/2018, 11/2019, 03/01/2020, 03/01/2022
AUTHOR(S): Visurraga Enterprises LLC, et al.	WEBSITE UPDATED: 03/01/2024
APPLIES TO: All Sites	APPROVED BY: Visurraga Enterprises LLC

References:

1. Institute for Clinical Systems Improvement. Healthcare Guideline: Perioperative (2020)

2. ACC/AHA Guideline: Perioperative Cardiac Evaluation and management of patients Undergoing non-cardiac surgery (2014).

3. Wanderer, P., Rathmell, J.P. (2017). Anesthesiology. Infographics in Anesthesiology, 126(21). doi:http://doi.org/10.1097/ALN.00000000001477

4. Lim, S., Rogers, L. K., Tessler, O., Mundinger, G. S., Rogers, C., & Lau, F. H. (2018). Phentermine: A Systematic Review for Plastic and Reconstructive Surgeons. Annals of plastic surgery, 81(4), 503-507.

5. Johns Hopkins Medicine (2019). Center for Perioperative Optimization. Preoperative Road

6. Girish P. Joshi, M.B.B.S., M.D., Basem B. Abdelmalak, M.D., Wade A. Weigel, M.D., Sulpicio G. Soriano, M.D., Monica W. Harbell, M.D., Catherine I. Kuo, M.D., Paul A. Stricker, M.D., Karen B. Domino, M.D., M.P.H., American Society of Anesthesiologists (ASA) Task Force on Preoperative Fasting (2023). American Society of Anesthesiologists Consensus-Based Guidance on Preoperative Management of Patients (Adults and Children) on Glucagon-Like Peptide-1 (GLP-1) Receptor Agonists.